

PUBLICATIONS ON THE KENTISH PLOVER IN PORTUGAL AND SPAIN (excluding doctoral thesis, atlas, faunistic studies and unpublished reports)

Compilation: Juan Aguilar Amat

- Alberto LJ, Velasco T (1988) Limícolas invernantes en España. In: Tellería JL (Ed). Invernada de aves en la Península Ibérica, pp 71-78. Monografías de la SEO, no 1. Madrid
- Almalki M, Kupán K, Carmona-Isunza MC, López P, Veiga A, Kostolányi A, Székely T, Küpper C (2017). Morphological differentiation among Kentish plover *Charadrius alexandrinus* populations in Macaronesia. *Ardeola* 64: 3-16.
- Alves JA, Lourenço PM, Dias MP, Antunes L, Catry T, Costa H, Fernandes P, Ginja B, Jara J, Martins R, Moniz F, Pardal S, Pereira T, Rainho MJ, Rocha A, Rodrigues JC, Rolo M (2012) Monitoring waterbird populations on the Tejo, Sado and Guadiana estuaries, Portugal: 2011 report. *Anuário Ornitológico* 9: 66-87.
- Amat JA (1993) Status of the Kentish plover in Spain. *Wader Study Group Kentish Plover Project Newsletter* 2: 2-4.
- Amat JA (1998) Mixed clutches in shorebird nests: why are they so uncommon? *Wader Study Group Bulletin* 85: 55-59.
- Amat JA (1999) Foot losses of metal banded snowy plovers. *Journal of Field Ornithology* 70: 555-557
- Amat JA (2005) Does the expression of a male plumage trait in female Kentish plovers (*Charadrius alexandrinus*) signal individual quality? *Journal of Ornithology* 146: 287-290.
- Amat JA, Fraga RM, Arroyo GM (1999) Reuse of nesting scrapes by Kentish plovers. *Condor* 101: 157-159.
- Amat JA, Fraga RM, Arroyo GM (1999) Replacement clutches by Kentish plovers. *Condor* 101: 746-751.
- Amat JA, Fraga RM, Arroyo GM (1999) Brood desertion and polygamous breeding in the Kentish plover *Charadrius alexandrinus*. *Ibis* 141: 596-607.
- Amat JA, Fraga RM, Arroyo GM (2001) Variations in body condition and egg characteristics of female Kentish plovers *Charadrius alexandrinus*. *Ardea* 89: 293-299.
- Amat JA, Fraga RM, Arroyo GM (2001) Intraclutch egg-size variation and offspring survival in the Kentish plover *Charadrius alexandrinus*. *Ibis* 143: 17-23.
- Amat JA, Masero JA (2004) Predation risk on incubating adults constrains the choice of thermally favourable nest sites in a plover. *Animal Behaviour* 67: 293-300.
- Amat JA, Masero JA (2004) How Kentish plovers, *Charadrius alexandrinus*, cope with heat stress during incubation. *Behavioral Ecology and Sociobiology* 56: 26-33.
- Amat JA, Masero JA (2007) The functions of belly-soaking in Kentish plovers *Charadrius alexandrinus*. *Ibis* 149: 91-97.

Socios / Parceiros:

Iberalex

- Amat JA, Masero JA (2009) Belly-soaking: a behavioural solution to reduce excess body heat in the Kentish plover *Charadrius alexandrinus*. *Journal of Ethology* 27:507-510.
- Amat JA, R. Monsa R, Masero JA (2012) Dual function of egg-covering in the Kentish plover *Charadrius alexandrinus*. *Behaviour* 149: 881-895.
- Amat JA, Visser GH, Pérez-Hurtado A, Arroyo GM (2000) Brood desertion by female shorebirds: a test of the differential parental capacity hypothesis on Kentish plovers. *Proceedings of the Royal Society of London B* 267: 2171-2176.
- Argüelles-Ticó A, Küpper C, Kelsh RN, Kosztolányi A, Székely T, van Dijk R (2016) Geographic variation in breeding system and environment predicts melanin-based plumage ornamentation of male and female Kentish plovers. *Behavioral Ecology and Sociobiology* 70: 49–60.
- Ballesteros T, Santaefemia FX (1990) Recapturas de chorlitejo patinegro (*Charadrius alexandrinus*) en el delta del Llobregat (Barcelona, NE Spain). *Butlletí del Grup Català d'Anellament* 7: 43-44.
- Ballesteros T, Torres I (1993) Incidencia de la predación sobre el fracaso de las puestas de chorlitejo patinegro *Charadrius alexandrinus* en el delta del Llobregat. *Butlletí del Grup Català d'Anellament* 10: 59-61.
- Barbosa A (1993). Ringing recoveries of Kentish plovers in Spain. *Wader Study Group Kentish Plover Project Newsletter* 2: 5-6.
- Barbosa A (1995) Foraging strategies and their influence on scanning and flocking behaviour of waders. *Journal of Avian Biology* 26: 182-186.
- Barbosa A (1997) Las aves limícolas en España. Colección Técnica, Organismo Autónomo de Parques Nacionales, Madrid.
- Barbosa A, Moreno E (1999). Evolution of foraging strategies in shorebirds: an ecomorphological approach. *Auk* 116: 712-725.
- Barbosa A, Moreno E (1999). Hindlimb morphology and locomotion performance in waders. An evolutionary approach. *Biological Journal of the Linnean Society* 67: 313-330.
- Batty L (1992) The wader communities of a saline and an intertidal site on the Ria Formosa, Portugal. *Wader Study Group Bulletin* 66: 66-72.
- Cabral JA, Pardal MÂ, Lopes RJ, Múrias T, Marques JC (1999) The impact of macroalgal blooms on the use of the intertidal area and feeding behaviour of waders (*Charadrii*) in the Mondego estuary (west Portugal). *Acta Oecologica* 20: 417-427.
- Castro M, Masero JA, Megina C, Amat JA, Pérez-Hurtado A (2008) Energy and macronutrient assimilation efficiencies of snowy plover (*Charadrius alexandrinus*) feeding on the polychaete ragworm (*Nereis diversicolor*). *Auk* 125: 368-373.
- Castro M, Masero JA, Pérez-Hurtado A, Amat JA, Megina C (2009) Sex-related seasonal differences in the foraging strategy of snowy plovers. *Condor* 111: 624-632
- Correia E, Granadeiro JP, Santos B, Regalla A, Mata VA, Catry T (2023) Trophic ecology of a migratory shorebird community at a globally important non-breeding site: combining DNA metabarcoding and conventional techniques. *Marine Ecology Progress Series* 705: 127–144.

Socios / Parceiros:

Iberalex

- Covelo P, Romero L, Mariscal P, Domínguez C, Marcuño P, Abad E, Valeiras J (2012) Voluntariado e custodia da nidificación da píllara das dunas (*Charadrius alexandrinus*) en Nigrán (Pontevedra) en 2012. Aplicación do "Plan de Conservación da Píllara das Dunas en Galicia". Memoria 2012. Sociedade Galega de Historia Natural (SGHN) e Coordinadora para o Estudo dos Mamíferos Mariños (CEMMA). 21pp.
- De Souza JA, Caeiro ML, Rosende F, Monteagudo A, Fafián, JM (1999) Estacionamientos, estrutura y patrones de residencia de la población invernante del chorlitejo patinegro (*Charadrius alexandrinus*) en Galicia: un análisis preliminar. *Chioglossa* 1: 23-45.
- De Souza JA, Domínguez J (1989). Efectivos y distribución del chorlitejo patinegro (*Charadrius alexandrinus*) en Galicia. *Ecología* 3: 305-311.
- De Souza JA, Fafián JM, Caeiro ML, Velasco J, Monteagudo A (1995) Situación actual del chorlitejo patinegro (*Charadrius alexandrinus*) en Galicia: población nidificante y primeros datos sobre productividad. In: Munilla I, Mouriño J (Eds). *Actas do II Congreso Galego de Ornitología*, pp. 95-113. Servicio de Publicacións e Intercambio Científico, Universidade de Santiago de Compostela, Santiago de Compostela.
- Dias MP, Peste F, Granadeiro JP, Palmeirim JM (2008) Does traditional shellfishing affect foraging by waders? The case of the Tagus estuary (Portugal). *Acta Oecologica* 33: 188-196.
- Domínguez J (1990) Distribution of estuarine waders wintering in the Iberian peninsula in 1978-1982. *Wader Study Group Bulletin* 59: 25-28.
- Domínguez J, Bárcena F, Souza JA, Villarino A (1987) Breeding waders in Galicia, north-west Spain. *Wader Study Group Bulletin* 50: 28-29.
- Domínguez J, Vidal M (2003) Influencia del investigador en el éxito reproductivo del chorlitejo patinegro *Charadrius alexandrinus*. *Ardeola* 50: 15-19.
- Domínguez J, Vidal M (2007). Proxecto Píllara. Potenciación e Estudo da Poboación reproductora da Píllara das dunas (*Charadrius alexandrinus*) nas praias de Galicia tras a catástrofe do Prestige. Fundación Arao. Santiago de Compostela.
- Domínguez J, Vidal M (2008) Plan de Conservación del Chorlitejo patinegro (*Charadrius alexandrinus*) en Galicia. Consellería de Medio Ambiente e Desenvolvemento Sostible. Santiago de Compostela.
- Domínguez J, Vidal M (2009) Factors affecting plumage oiling levels in a Kentish plover *Charadrius alexandrinus* population after a major oil spill. *Acta Ornithologica* 44: 119–126.
- D'Urban Jackson J, dos Remedios N, Maher KH, Zefania S, Haig S, Oyler-McCance S, Blomqvist D, Burke T, Bruford MW, Székely T, Küpper C (2017) Polygamy slows down population divergence in shorebirds. *Evolution* 71: 1313–1326.
- Ferreira-Rodríguez N, Pombal MA (2018) Predation pressure on the hatching of the Kentish plover (*Charadrius alexandrinus*) in clutch protection projects: a case study in north Portugal. *Wildlife Research* 45: 55–63.
- Figuerola J, Cerdà F. (1998). Evolució i conservació de la població de corriol camanegre (*Charadrius alexandrinus*) del delta del Llobregat. *Spartina* 3: 161-169.

Socios / Parceiros:

Iberalex

- Figuerola J, Velar R, Bertolero A, Cerdá F (1996) Abwesenheit von Haematozoa bei einer Brutpopulation des Seeregenpfeifers *Charadrius alexandrinus* in Nordspanien. Journal für Ornithologie 137: 523-525.
- Fonseca VG, Grade N, Cancela da Fonseca, L (2004) Patterns of association and habitat use by migrating shorebirds on intertidal mudflats and saltworks on the Tavira Estuary, Ria Formosa, southern Portugal. Wader Study Group Bulletin 105: 50–55.
- Fonseca VG, Grade N, Cancela da Fonseca, L (2005) Waterbird breeding on salinas in Ria Formosa, southern Portugal. Wader Study Group Bulletin 106: 58-59.
- Forner E, Roig-Munar FX, Prades R (2021) Nedar a contracorrent: evolució poblacional del corriol camanegre, *Charadrius alexandrinus*, (1989-2019) a la primera línia de platja de Castelló de la Plana (País Valencià). Bolletí de la Societat d'Història Natural de les Balears 64: 61-82.
- Forteza V, Roig-Munar FX, Colomar JL, Ramis M, Galludo M, Castelló M, Valverde N (2020) Noves zones de nidificació de *Charadrius alexandrinus* (picaplatges camanegre), al parc natural de ses Salines d'Eivissa i Formentera (illes Balears): efectes i conseqüències de la declaració de l'estat d'alarma per la COVID-19 i mesures de gestió. Nemus 10: 71-85.
- Fraga RM, Amat JA (1996) Breeding biology of a Kentish plover (*Charadrius alexandrinus*) population in an inland saline lake. Ardeola 43: 69-85.
- García López de Haro E (2022) Crecimiento, desarrollo y supervivencia de los pollos de chorlito patinegro *Charadrius alexandrinus* en ambientes hipersalinos de la Bahía de Cádiz, Trabajo de Fin de Máster, Universidad de Cádiz.
- Garcias PJ, Tavecchia, G (2018) Apparent survival and long-term population growth rate of the Kentish plover *Charadrius alexandrinus* in Mallorca, Balearic Archipelago, Spain. Ardeola 65: 59-68.
- Gomes JT (2022) Waterbirds as sentinelas of microplastic and plastic additive pollution in coastal systems in Southern Portugal. Dissertação do Mestrado em Ecologia, Universidade de Coimbra.
- Gómez J, Liñán-Cembrano G, Castro M, Pérez-Hurtado A, Ramo C, Amat JA (2019) The protoporphyrin content of Kentish plover *Charadrius alexandrinus* eggshells is better predicted by the fractal dimension of spottiness than by colour. Journal of Ornithology 160: 409-415 (2019)
- Gómez J, Liñán-Cembrano G, Ramo C, Castro M, Pérez-Hurtado A, Amat JA (2019) Does the use of nest materials in a ground-nesting bird result from a compromise between the risk of egg overheating and camouflage? Biology Open 8: bio042648
- Gómez J, Pereira AI, Pérez-Hurtado A, Castro M, Ramo C, Amat JA (2016) A trade-off between overheating and camouflage on shorebird eggshell colouration. Journal of Avian Biology 47: 346-353.
- Gómez J, Ramo C, Stevens M, Liñán-Cembrano G, Rendón MA, Troscianko JT, Amat JA (2018) Latitudinal variation in biophysical characteristics of avian eggshells to cope with differential effects of solar radiation. Ecology and Evolution 8: 8019-8029.

Socios / Parceiros:

Iberalex

- Gómez J, Ramo C, Troscianko JT, Stevens M, Castro M, Pérez-Hurtado A, Liñán-Cembrano G, Amat JA (2018) Individual egg camouflage is influenced by microhabitat selection and use of nest materials in ground-nesting birds. *Behavioral Ecology and Sociobiology* 72: 142.
- Gómez J, Ramo C, Canales JA, García IM, Castro M, Pérez-Hurtado A, Amat JA (2020) A comparative assessment of pigment extraction from avian eggshells using two analytical protocols. *Ardeola* 67: 85-92.
- Gómez-Serrano MÁ (2021) Four-legged foes: dogs disturb nesting plovers more than people do on tourist beaches. *Ibis* 163: 338-352.
- Gómez-Serrano MÁ, López-López P (2014) Nest site selection by Kentish plover suggests a trade-off between nest-crypsis and predator detection strategies. *PLoS ONE* 9: e107121.
- Gomez-Serrano MÁ, López-López P (2017) Deceiving predators: linking distraction behavior with nest survival in a ground-nesting bird. *Behavioral Ecology* 28: 260–269.
- Gómez-Serrano MA, Prades R, Hernández-Navarro VJ (1994) Efectivos y distribución de las aves limícolas nidificantes en Castellón. *Evolución de las poblaciones. Actas de las XII Jornadas Ornitológicas, SEO-BirdLife*. 87-97.
- Gómez-Serrano MÁ, Prades R (1997) Conservación de las poblaciones nidificantes de aves limícolas en Castellón. In: Manrique J, Sánchez A, Suárez F, Yanes M (Eds.) *Actas de las XII Jornadas Ornitológicas Españolas*, pp. 79-85. Instituto de Estudios Almerienses, Almería.
- Gonçalves MSS, Gil-Delgado JA, Gosálvez RU, López-Iborra GM, Ponz A, Velasco A (2016) Spatial synchrony of wader populations in inland lakes of the Iberian Peninsula. *Ecological Research* 31: 947–956.
- Granadeiro JP, Andrade J, Palmeirim JM (2004) Modelling the distribution of shorebirds in estuarine areas using generalised additive models. *Journal of Sea Research* 52: 227-240.
- Granadeiro JP, Santos CD, Dias MP, Palmeirim JM (2007) Environmental factors drive habitat partitioning in birds feeding in intertidal flats: implications for conservation. *Hydrobiologia* 587: 291–302.
- Guinart E, Burgas A, Ramos S, Martí-Aledo, J, Roig-Munar, FX (2021) La gestió dunar en espais naturals protegits del litoral de Girona i el seu efecte en la població del corriol camanegre (*Charadrius alexandrinus*) durant el període 2012- 2020. *Bolletí de la Societat d'Història Natural de les Balears* 64: 91-100.
- Halimubieke N, Krisztina Kupán K, Valdebenito JO, Kubelka V, Carmona-Isunza MC, Burgas D, Catlin D, St Clair JJH, Cohen J, Figuerola J, Yasué M, Johnson M, Mencarelli M, Cruz-López M, Stantial M, Weston MA, Lloyd P, Que P, Montalvo T, Bansal U, McDonald GC, Liu Y, Kosztolányi A, Székely T (2020) Successful breeding predicts divorce in plovers. *Scientific Reports* 10: 15576.
- Hortas F (1995). Population tendencies and distribution of the Kentish plover in industrial salines of Cadiz Bay, southwest of Spain. *Wader Study Group Kentish Plover Project Newsletter* 4: 9.

Socios / Parceiros:

Iberalex

- Hortas F, Pérez-Hurtado A, Robledano F, Álvarez Laó C, Salvadores R (2008) Population estimates of waders on Spanish non-estuarine coasts. *International Wader Studies* 18: 65-74.
- Küpper C, Edwards EW, Kosztolányi A, Alrashidi M, Burke T, Herrmann P, Argüelles-Tico A, Amat JA, Amezian M, Rocha A, Hötker H, Ivanov A, Chernicko J, Székey T (2012) High gene flow on a continental scale in the polyandrous Kentish plover *Charadrius alexandrinus*. *Molecular Ecology* 21: 5864-5879.
- Lecoq M, Lourenço PM, Catry P, Andrade J, Granadeiro JP (2013). Wintering waders on the Portuguese mainland non-estuarine coast: results of the 2009–2011 survey. *Wader Study Group Bulletin* 120: 66-70.
- Lopes RJ, Múrias T, Cabral JA, Marques JC (2005) A ten year study of variation, trends and seasonality of a shorebird community in the Mondego Estuary, Portugal. *Waterbirds*, 28: 8-18.
- Lorenzo JA (1993) A case of three clutches in the same nest by the same pair of Kentish plover *Charadrius alexandrinus*. *Wader Study Group Bulletin* 71: 25-26.
- Lorenzo JA (1993) Descripción de la comunidad de aves limícolas de El Médano (Tenerife, Islas Canarias) durante un ciclo anual. *Ardeola* 40: 13-19.
- Lorenzo JA, Emmerson KW (1995) Recent information on the distribution and status of the breeding population of Kentish plover *Charadrius alexandrinus* in the Canary Islands. *Wader Study Group Bulletin* 76: 43-46.
- Lorenzo JA, Emmerson KW (1996) Summering coastal waders on Fuerteventura, Canary Islands, Spain. *Wader Study Group Bulletin* 79: 87-90.
- Lorenzo JA, González J (1993) Datos sobre la biología del chorlitejo patinegro (*Charadrius alexandrinus*) en la última población nidificante en la isla de Tenerife con vistas a su futura protección y conservación. *Alytes* 6: 199-219.
- Lorenzo JA, González J (1995). Las aves limícolas de Canarias: breve sinopsis y estado actual. *Airo* 6: 7-14.
- Lourenço PM (2006) Seasonal abundance of aquatic birds at Óbidos Lagoon. *Airo* 16: 23-29
- Lourenço PM, Alonso H, Alves JA, Carvalho AT, Catry T, Costa H, Costa JS, Dias MP, Encarnação V, Fernandes P, Leal AI, Martins RC, Moniz F, Pardal S, Rocha A, Santos CD (2018) Monitoring waterbird populations in the Tejo estuary, Portugal: report for the decade 2007-2016. *Airo* 25: 3-31.
- Lourenço PM, Catry P, Lecoq M, Ramírez I, Granadeiro JP (2013) Role of disturbance, geology and other environmental factors in determining abundance and diversity in coastal avian communities during winter. *Marine Ecology Progress Series* 479: 223–234.
- Lourenço PM, Silva A, Santos CD, Miranda AC, Granadeiro JP, Palmeirim JD (2008) The energetic importance of night foraging for waders wintering in a temperate estuary. *Acta Oecologica* 34: 122-129.
- Maher KH, Eberhart-Phillips LJ, Kosztolányi A, dos Remedios N, Carmona-Isunza MC, Cruz-López M, Zefania S, St Clair JJH, Alrashidi M, Weston MA, Serrano-Meneses MA, Krüger O, Hoffman JI, Székely T, Burke T, Küpper C (2017) High fidelity: extra-pair

Socios / Parceiros:

Iberalex

fertilisations in eight *Charadrius* plover species are not associated with parental relatedness or social mating system. *Journal of Avian Biology* 48: 910–920.

- Martín A, Mulet T (2012) Censo de la población reproductora de chorlitojo patinegro (*Charadrius alexandrinus*) en el cordón dunar de Gandía a Oliva (Valencia). *El Serenet* 8: 25-32.
- Martín B, Delgado S, de la Cruz A, Tirado S, Ferrer M (2015) Effects of human presence on the long-term trends of migrant and resident shorebirds: evidence of local population declines. *Animal Conservation* 18: 73–81
- Martínez Vilalta A (1985). Breeding waders of the Iberian Peninsula. *Wader Study Group Bulletin* 45: 35-36.
- Martínez-Vilalta A (1985). Descripción de la comunidad de limícolas invernante en el Delta del Ebro. *Doñana Acta Vertebrata* 12: 211-229.
- Masero JA, Pérez-Hurtado A, Castro M, Arroyo GM (2000). Complementary use of intertidal mudflats and adjacent salinas by foraging waders. *Ardea* 88: 177-191.
- Montalvo T, Figuerola J (2006) The distribution and conservation of the Kentish plover *Charadrius alexandrinus* in Catalonia. *Revista Catalana d'Ornitologia* 22:1-8.
- Munn PW (1948) The nesting of Kentish and little ringed plovers in Mallorca. *Ibis* 90: 595.
- Múrias T, Cabral JA, Marques JC, Goss-Custard JD (1996) Short-term effects of intertidal macroalgal blooms on the macrohabitat selection and feeding behaviour of wading birds in the Mondego estuary (west Portugal). *Estuarine, Coastal and Shelf Science* 43: 677-688.
- Múrias T, Cabral JA, Lopes R, Marques JC (1997) Low-water use of the Mondego Estuary (West Portugal) by wader (*Charadrii*). *Ardeola* 44:79-91.
- Múrias T, Cabral JA, Lopes RJ, Marques JC, Goss-Custard J (2002). Use of traditional salines by waders in the Mondego estuary (Portugal): A conservation perspective. *Ardeola* 49: 223-240.
- Múrias T, Cabral JA, Lopes RJ, Pardal M, Marques JC, Goss-Custard J (2005) Competition for feeding in waders: a case study in an estuary of south temperate Europe (Mondego, Portugal). *Hydrobiologia* 544: 155–166.
- Múrias T, Gonçalves D, Lopes RJ (2017) Evaluating the impacts of a new railway on shorebirds: A case study in central Portugal (Aveiro Lagoon). In: Borda-de-Água L, Barrientos R, Beja P, Pereira HM (Eds) *Railway Ecology*, pp. 197-214. Springer, Cham.
- Novo Varela N (2022) Modelización do hábitat reproductor de *Charadrius alexandrinus* (Linnaeus, 1758) na costa de Galicia mediante imaxes de satélite. Trabajo de Fin de Mestrado, Universidade da Coruña.
- Ollé À, Martí-Aledo J (2008) Resum de la temporada nidificant 2008 del corriol camanegre *Charadrius alexandrinus* als Aiguamolls de l'Empordà. *Butlletí de l'Institut d'Estudis Altampordanesa per a la Defensa i Estudi de la Natura* 131: 27-29.
- Ollé À, Martí-Aledo J, Burgas A (2009) Últimes tendències poblacionals del corriol camanegre *Charadrius alexandrinus* a l'Empordà. *Annals de l'Institut d'Estudis Empordanesos* 40: 163-174.

Socios / Parceiros:

Iberalex

- Oltra Martínez C, Gómez Serrano MA (1993) Situación de la población nidificante de chorlitejo patinegro en la Comunidad Valenciana en 1993. Anuario Ornitológico de la Comunidad Valenciana 1993: 146-148.
- Oltra O, Gómez-Serrano MA (1993) Situation of breeding population of Kentish Plover (*Charadrius alexandrinus*) in Valencian Area (Spain) in 1993. First results. Wader Study Group Kentish Plover Project 3: 12-15.
- Oltra O, Gómez-Serrano MA (1997) Amenazas humanas sobre las poblaciones nidificantes de limícolas en ecosistemas litorales. In Las aves limícolas en España (pp. 175-200). Ministerio de Medio Ambiente, Colección Técnica.
- Orr N (1999) Egg-burying behaviour of Kentish plovers. British Birds 92: 365-372.
- Paracuellos M, Castro H, Nevado JC, Oña JA, Matamala, García L, Salas G (2002) Repercussions of the abandonment of Mediterranean salt pans on waterbird communities. Waterbirds 25: 492-498.
- Pedro P, Ramos JA (2009) Diet and prey selection of shorebirds on salt pans in the Mondego estuary, western Portugal. Ardeola 56: 1-11
- Pérez-Hurtado A (1995) Ecología alimentaria de limícolas invernantes en la Bahía de Cádiz. Airo 6: 15-23.
- Pérez-Hurtado A, Goss-Custard JD, García F (1997) The diet of wintering waders in Cádiz Bay, southwest Spain. Bird Study 44: 45-52.
- Pérez-Hurtado A, Hortas F (1993). Actividad trófica de limícolas invernantes en salinas y cultivos piscícolas de la Bahía de Cádiz. Doñana Acta Vertebrata 20: 103-123.
- Pérez-Hurtado A, Hortas F, Ruiz F, Solís F (1993) Importancia de la Bahía de Cádiz para las poblaciones de limícolas invernantes e influencia de las transformaciones humanas. Ardeola 40: 133-142.
- Polo M (2014) Un caso de parasitismo reproductor entre chorlitejo chico (*Charadrius dubius*) y chorlitejo patinegro (*Charadrius alexandrinus*). El Serenet 9: 7-11.
- Queral I, Sánchez FJ, Prades R (1993) Ecología del *Charadrius alexandrinus* en un tramo protegido del litoral castellonense (Comunidad Valenciana). Alytes 6: 221-229.
- Quesada J, Pàmies E, Oliver C, MacGregor-Fors I (2022) Hunter sparrows: a predation record of a house sparrow on Kentish plover nestlings. Revista Catalana d'Ornitologia 38: 37-42.
- Ramos J, Pinto P, Pintassilgo P, Resende A, Cancela da Fonseca L (2022) Activating an artisanal salt pan: tourism crowding in or waterbirds crowding out? International Journal of Culture, Tourism and Hospitality Research 16: 294-305.
- Robledano F (1995) Ecología de humedales y conservación de limícolas en el SE de España: experiencias de investigación aplicables a la gestión. Airo 6: 39-56.
- Rocha AD, Fonseca D, Masero JA, Ramos JA (2016) Coastal salt pans are a good alternative breeding habitat for Kentish plover *Charadrius alexandrinus* when umbrella species are present. Journal of Avian Biology 47: 824–833.

Socios / Parceiros:

Iberalex

- Rosa S, Encarnação AL, Granadeiro JP, Palmeirim JM (2006) High water roost selection by waders: maximizing feeding opportunities or avoiding predation? *Ibis* 148: 88-97.
- Rubio JC (1986) Estudio de la comunidad de limícolas de las Marismas del Odiel (Huelva). *Oxyura* 3: 97-132.
- Rufino R, Araújo A (1987) Seasonal variations in wader numbers and distribution at the Ria de Faro. *Wader Study Group Bulletin* 51: 48-53.
- Rufino R, Araújo A, Pina JP, Miranda PS (1984) The use of salinas by waders in the Algarve, south Portugal. *Wader Study Group Bulletin* 42: 41-42
- Rufino R, Neves R (1992) The effects on wader populations of the conversion of the salinas into fish farms. *IWRB Special Publication No. 20. Proceedings of the Symposium Managing Mediterranean Wetlands and their Birds for the Year 2000 and Beyond*: 177-182.
- Santaefemia FJ, Ballesteros T, García J, Puig M (1990) Características de la población nidificante del chorlitejo patinegro (*Charadrius alexandrinus*) en el Delta del Llobregat. *Butlletí Parc Natural Delta de l'Ebre* 5: 31-34.
- Székely T (1996) Brood desertion in Kentish plover *Charadrius alexandrinus*: an experimental tests of parental quality and remating opportunities. *Ibis* 138: 749-755.
- Tavares PC, McGill R, Newton J, Pereira E, Duarte A, Furness RW (2009) Relationships between carbon sources, trophic level and mercury exposure in generalist shorebirds revealed by stable isotope ratios in chicks. *Waterbirds* 32: 311-321.
- Tejera G, Amat JA, Rodríguez B, Arizaga J (2022) Apparent survival, reproduction and population growth estimation of a Kentish plover population in the Canary Islands. *European Journal of Wildlife Research* 68: 52.
- Tejera G, Rodríguez B (2014) Quantifying the importance for waterbirds of an urban rocky coastal site in Lanzarote, Canary Islands. *Wader Study Group Bull.* 121: 1-8.
- Toral GM, Figuerola J (2012) Nest success of black-winged Stilt *Himantopus himantopus* and Kentish plover *Charadrius alexandrinus* in rice fields, southwest Spain. *Ardea* 100: 29-36.
- Torres I, Ballesteros T (1994) Variabilidad en el tamaño del huevo del chorlitejo patinegro (*Charadrius alexandrinus*) en el Delta del Llobregat. *Butlletí del Grup Català d'Anellament* 11: 89-92.
- Valdebenito JO, Martínez-de la Puente J, Castro M, Pérez-Hurtado A, Tejera G, Székely T, Halimubieke N, Schroeder J, Figuerola J (2020) Association of insularity and body condition to cloacal bacteria prevalence in a small shorebird. *PLoS ONE* 15: e0237369.
- Vázquez A, Vázquez J (1998) Migració prenupcial de limícoles als Salats i Muntanyans del Tarragonès. *Butlletí del Grup Català d'Anellament* 15: 33-37.
- Velasco T (1996) La migración postnupcial de limícolas en una playa malagueña durante el mes de agosto. *Butlletí del Grup Català d'Anellament* 13: 15-19.
- Velasco T, Alberto LJ (1993) Numbers, main localities and distribution maps of waders wintering in Spain. *Wader Study Group Bulletin* 70: 33-41.

Socios / Parceiros:

Iberalex

- Velasco T, Alberto LJ (1994) Wintering of Kentish plover *Charadrius alexandrinus* in Spain: numbers, distribution and habitat selection. Wader Study Group Bulletin 73: 54-55.
- Vidal M, Domínguez J (2013) Long-term population trends of breeding Kentish plovers *Charadrius alexandrinus* in north-west Spain under the effects of a major oil spill. Bird Conservation International 23: 386–397.
- Vidal M, Domínguez J (2015) Did the Prestige oil spill compromise bird reproductive performance? Evidences from long-term data on the Kentish plover (*Charadrius alexandrinus*) in NW Iberian Peninsula. Biological Conservation 191: 178-184.
- Vidal M, Domínguez J, Luís A (2011). Spatial and temporal patterns of polycyclic aromatic hydrocarbons (PAHs) in eggs of a coastal bird from northwestern Iberia after a major oil spill. Science of the Total Environment 409: 2668-2673.
- Vidal M, García-Miranda Ó, Domínguez J (2021) Sex-related and spatial variation in trace elements in feathers of the Kentish plover (*Charadrius alexandrinus*) from different coastal areas of the Iberian Peninsula. Science of the Total Environment 766: 144628.
- Vidal M, Hernández MA, Luis A, Domínguez J (2015) Lack of genetic structure in Kentish plover *Charadrius alexandrinus* from the Iberian coast. What's wrong with the endangered north-western population? Bird Conservation International 25: 153 –165.
- Vincze O, Kosztolányi A, Barta Z, Küpper C, Alrashidi M, Amat JA, Argüelles Ticó A, Burns F, Cavitt J, Conway WC, Cruz-López M, Desucre-Medrano AE, dos Remedios N, Figuerola J, Galindo-Espinosa JD, García-Peña GE, Gómez del Angel S, Gratto-Trevor SC, Jönsson P, Lloyd P, Montalvo T, J. Parra JE, Pruner R, Que P, Liu Y, Saalfeld ST, Schulz R, Serra L, St Clair JJH, Stenzel LE, Weston MA, Yasué M, Zefania S, Székely T (2017) Parental cooperation in a changing climate: fluctuating environments predict shifts in care division. Global Ecology and Biogeography 26: 347-358.
- Vincze, O., T. Székely, C. Küpper, M. AlRashidi, J. A. Amat, A. Argüelles Ticó, D. Burgas, T. Burke, J. Cavitt, J. Figuerola, M. Shobrak, T. Montalvo y A. Kosztolányi A (2013) Local environment but not genetic differentiation influences biparental care in ten plover populations. PLoS ONE 8: e60998.

Socios / Parceiros: